

Major Article

Phage Therapy as an Approach to Control *Salmonella enterica* serotype Enteritidis Infection in Mice

Mohammad Mehdi Soltan Dallal^{[1],[2]}, **Farhad Nikkhahi**^{[3],[4]}, **Mahmoud Alimohammadi**^[5], **Masoumeh Douraghi**^{[2],[3]}, **Zahra Rajabi**^[2], **Abbas Rahimi Foroushani**^[6], **Akram Azimi**^{[3],[4]} and **Fatemeh Fardsanei**^[3]

[1]. Division of Food Microbiology, Department of Pathobiology, School of Public Health (TUMS), Tehran, Iran.

[2]. Food Microbiology Research Center, Tehran University of Medical Sciences (TUMS), Tehran, Iran.

[3]. Division of Medical Bacteriology, Department of Pathobiology, School of Public Health (TUMS), Tehran, Iran.

[4]. Medical Microbiology Research Center, Qazvin University of Medical Science, Qazvin, Iran.

[5]. Environmental Health Engineering Department, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran.

[6]. Department of Epidemiology and Biostatistics, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran.

Abstract

Introduction: *Salmonella enterica* serotype Enteritidis (*S. Enteritidis*) is a cause of food-borne human illness. Given the prevalence of antibiotic resistance of *Salmonella* Enteritidis and the lack of antibiotic efficacy in future years, its replacement with other agents is necessary. One of the most useful agents is bacteriophages. **Methods:** *S. Enteritidis* was identified using a multiplex polymerase chain reaction assay. The effective bacteriophages were isolated from hospital wastewater samples. The effects of the bacteriophages were evaluated both *in vitro* and *in vivo*. **Results:** The phage SE20 belonged to the Podoviridae family, and the genome size was 40 kb. The evaluation of phage SE20 at variable pH ranges showed its susceptibility to pH < 3 and pH > 12. The animal model showed that mice infected with *S. Enteritidis* developed hepatomegaly and splenomegaly, but did not experience gastrointestinal complications after receiving the bacteriophages. **Conclusions:** The results of this study suggest that phage SE20 is a promising candidate for controlling salmonellosis caused by *Salmonella* Enteritidis.

Keywords: *Salmonella enterica* serotype Enteritidis. Phage SE20. Pulsed-field gel electrophoresis. Mice. Sewage.